IN THE KNOW:

AND OTHER **OPIATES**

Types of Opioids

Illegal

 Heroin Desomorphine

(krokodil)

Acetyl Fentanyl

Prescription

- Morphine
- Codeine
- Oxycodone
- Hydrocodone
- Fentanyl
- Methadone

Signs of Abuse

- Excessive sleeping
- · Difficulty moving
- Weight gain/loss
- Poor hygiene
- · Hanging out with new groups of people
- · Track/needle marks and wearing long sleeves to hide them
 - · Borrowing money with no reason aiven

Opioid Facts

lunhuhuh **P**unhuh

Every year, about 150,000 people in the U.S. try heroin for the first time.

About **2,200,000** peopl in the U.S. are addicted to opioids.

Addicts will spend \$150 to \$200 per day on their habit.



Effects:

Long-Term

Effects:



- · Poor coordination
- Slowed breathing Lowered heart rat
- Coma
- Brain damage • Endorphin deficit
- Hormonal imbalance **Emotional difficulties**
- Osteoporosis



Addiction & Dependency



Addiction can happen after just one use Addicts will:

Crave the drug.

Be compelled to use the drug, even if they know there will be negative consequences.

Not be able to control their drug use.

Build up a tolerance to the drug.

Overdose & Withdrawal

Overdose is a real danger. Take too much and the body literally shuts down:

- Breathing and heart rate slow, maybe even stop.
- Lips and nails will turn blue because of insufficient oxygen in the blood.
- Seizure or muscle spasms may occur.
- Vomiting and choking may also happen.



Every 18 minutes in the U.S. someone dies from an accidental overdose.

Withdrawal & Detox Opioid withdrawal is

painful and can last for up to two weeks. Symptoms include:

- Nausea
- Muscle and bone pain
- Excessive sweating
- Anxiety and agitation
- Diarrhea
- Stomach cramps
- Vomiting

90% of opioid addicts will relapse in the first year.

Getting Help



Overcoming opioid addiction is difficult – but it's not impossible! Residential, inpatient, and outpatient services and support grou can provide the reinforcement you need in order to quit.

Avoid Addiction - Avoid Opioids