APPROXIMATELY 32,537 PEOPLE DIED FROM AN OVERDOSE INVOLVING STIMULANTS LIKE AS METH.

METH HEART FAILURE IS ON THE RISE WORLDWIDE, AND AFFECTS A WIDE RANGE OF RACIAL/ETHNIC AND SOCIOECONOMIC GROUPS.

ABOUT 2.5 MILLION PEOPLE AGED 12 OR OLDER REPORTED USING METH IN THE PAST 12 MONTHS.

ALMOST 1 IN 5 (18%) DEVELOP HEART FAILURE WITHIN 12 MONTHS OF METH USE

USING METH DURING PREGNANCY INCREASES THE RISK OF LOW BIRTH WEIGHT AND HEART OR BRAIN ABNORMALITIES

OVER A FOUR YEAR PERIOD, DEATHS FROM METH OVERDOSES INCREASED 170%

AN EXAMINATION OF THE MOUTHS OF METHAMPHETAMINE USERS SHOWED: 96% OF METH USERS HAD CAVITIES AND 31% HAD SIX OR MORE MISSING TEETH.

HEART DISEASE REPRESENTS THE SECOND-LEADING CAUSE OF DEATH AMONG THOSE WHO USE METHAMPHETAMINE FOLLOWING ONLY ACCIDENTAL OVERDOSE

PREVIEW COPY